

## MP.008.MPC Home Apnea Monitoring

Maryland Physicians Care considers **Home Apnea Monitors** medically necessary when they are equipped with an event recorder and are indicated for a limited period of time for infants 12 months of age or younger with *any* of the following indications:

- An infant who has experienced a Brief Unexplained Resolved Event (BRUE) <sup>(1, 3)</sup>
- Premature infants who are at high risk for recurrent episodes of apnea <sup>(2)</sup>
- Bradycardia to less than 80 beats per minute, and hypoxia (oxygen saturation below 90%) after discharge from the hospital <sup>(2)</sup>
- Infants who are technology dependent – tracheostomy, Continuous Positive Airway Pressure (CPAP), or mechanical ventilation <sup>(6)</sup>
- Infants with unstable airways <sup>(6)</sup>
- Infants with neurologic or metabolic disorders affecting respiratory control or rare medical conditions that affect regulation of breathing <sup>(1, 6)</sup>
- Chronic lung disease
- Infants with confirmed diagnosis of pertussis <sup>(7)</sup>
- Later siblings of infants who died of Sudden Infant Death Syndrome (SIDS) until the siblings are one month older than the age at which the earlier sibling died and they remain event free. <sup>(1)</sup>

The physician must establish a specific plan for periodic review and criteria for termination of the home monitor before initiating therapy. Parents require supportive care and education and need to be advised that home monitoring has never been demonstrated to reduce the rate of mortality caused by Sudden Infant Death Syndrome (SIDS). <sup>(4)</sup>

Infant apnea monitors are experimental and investigational for all other indications because their effectiveness for indications other than the ones listed above has not been established.

### Limitations

1. Home apnea monitors should be discontinued after infants are event-free (no episodes of apnea/bradycardia) for six weeks and post-conception age of 43 weeks.
2. The use of the apnea monitor is not indicated for the sole purpose of prevention of SIDS without a history of sibling SIDS.

## MP.008.MPC Home Apnea Monitoring

Policy Number: MP.008.MPC

Last Review Date: 02/20/2025

Effective Date: 03/01/2025

3. This policy will follow the capped rental period – see policy PA.010 Durable Medical Equipment and Corrective Appliances.

### Codes

#### CPT Codes

Code	Description
94774	Pediatric home apnea monitoring event recording including respiratory rate, pattern and heart rate per 30-day period of time includes monitor attachment, download of data, physician review, interpretation, and preparation of a report
94775	Pediatric home apnea monitoring event recording including respiratory rate, pattern and heart rate per 30-day period of time: includes monitor attachment only (includes hook-up, initiation of recording and disconnection)
94776	Pediatric home apnea monitoring event recording including respiratory rate, pattern and heart rate per 30-day period of time: monitoring, download of information, receipt of transmissions(s) and analyses by computer only
94777	Pediatric home apnea monitoring event recording including respiratory rate, pattern and heart rate per 30-day period of time: physician review, interpretation and preparation of report only

### References

1. Dick, A. Outcomes for the apparent life-threatening event infant. Health Science Reports. 2023 Feb; 6:e1152. Outcomes for the apparent life-threatening event infant  
<https://pubmed.ncbi.nlm.nih.gov/36938143/>
2. Eichenwald, EC. Apnea of Prematurity. Pediatrics. 2023; 137 (1): e20153757. Apnea of Prematurity | Pediatrics | American Academy of Pediatrics  
<https://pubmed.ncbi.nlm.nih.gov/26628729/>
3. Hall KL, Zalman B. Evaluation and management of apparent life-threatening events in children. Am Fam Physician. 2005 Jun; 71(12): 2301-2308.  
<https://www.aafp.org/pubs/afp/issues/2005/0615/p2301.html>
4. Moon RY, Carlin RF, Hand I. Evidence Base for 2022 Updated Recommendations for a Safe Infant Sleeping Environment to Reduce the Risk of Sleep-Related Infant Deaths. Paediatrics. 2022 Jul;150 (1):e2022057991.  
<https://pubmed.ncbi.nlm.nih.gov/35921639/>
5. Rocker JA, Bechtel KA. Pediatric Apnea. Last updated: July 18, 2021.  
<http://emedicine.medscape.com/article/800032-overview>

## MP.008.MPC Home Apnea Monitoring

Policy Number: MP.008.MPC

Last Review Date: 02/20/2025

Effective Date: 03/01/2025

6. Velumula P, Jani S, Kanike N, Chawla S. Monitoring of Infants Discharged Home with Medical Devices. *Pediatric Annals*. 2020 Nov; 49 (2). [Monitoring of Infants Discharged Home with Medical Devices | Pediatric Annals](https://pubmed.ncbi.nlm.nih.gov/32045488/)  
<https://pubmed.ncbi.nlm.nih.gov/32045488/>
7. Lee J, Robinson J and Spady D. Frequency of apnea, bradycardia, and desaturations following first diphtheria-tetanus-pertussis-inactivated polio-Haemophilus influenza type B immunization in hospitalized preterm infants. *BMC Pediatrics*. 2006; 6 (20).  
<https://bmcpediatr.biomedcentral.com/articles/10.1186/1471-2431-6-20>

### Disclaimer

Maryland Physicians Care medical payment and prior authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. The policies constitute only the reimbursement and coverage guidelines of Maryland Physicians Care and its affiliated managed care entities. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies.

Maryland Physicians Care reserves the right to review and update the medical payment and prior authorization guidelines in its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

These policies are the proprietary information of Maryland Physicians Care. Any sale, copying, or dissemination of said policies is prohibited.